

# FLOXY-SPRAY

Florfenicol 2,5%



MODERN ALTERNATIVE  
TO TRADITIONAL SPRAYS



# FLORFENICOL: SETTING A NEW STANDARD IN ANTIMICROBIAL SPRAY EFFECTIVENESS

## BROAD-SPECTRUM ACTIVITY


Gram<sup>+</sup> / Gram<sup>-</sup> bacteria,  
mycoplasmas

## ACTIVITY AGAINST GRAM- NEGATIVE BACTERIA

Including tetracycline-  
resistant strains

## LOW RISK OF RESISTANCE

Due to a different  
mechanism of action



## HIGH STABILITY

Effectiveness in  
purulent-necrotic lesions

## FLOXY-SPRAY VS. TRADITIONAL SPRAYS

|                                  | TETRACYCLINE SPRAYS                                     |                                 | FLOXY-SPRAY                       |
|----------------------------------|---------------------------------------------------------|---------------------------------|-----------------------------------|
| Active substance                 | Chlortetracycline                                       | Oxytetracycline                 | Florophenicol                     |
| Types of action                  | Bacteriostatic                                          | Bacteriostatic                  | Bacteriostatic                    |
| Effect on Gram-positive bacteria | High                                                    | High                            | High                              |
| Effect on Gram-negative bacteria | Often low, there is a risk of developing resistance     | Moderate, depends on the strain | High, including resistant strains |
| Resistance                       | High (TET genes), especially with long-term use         |                                 | Low                               |
| Stability in necrotic wounds     | Low — (organic materials, Ca/Mg ions reduce efficiency) |                                 | High                              |

**FLOXY-SPRAY** is a modern alternative with a broad spectrum of action, high efficacy against tetracycline-resistant bacteria, and stability in severe cases.

# MAXIMUM CONFIDENCE WITH EVERY PRESS



## OPTIMIZED FORMULATION

- No crystallization of contents
- Uniformity during storage
- Safe food coloring

## HIGH-QUALITY SPRAY-BOTTLE

- Uniform spray
- High sealing
- Ergonomic design

## LONG-ACTING

- Reliable skin contact
- Deep penetration into the infection site
- Protection from recontamination



# INDICATIONS AND ADMINISTRATION


| INDICATIONS FOR USE                                                                               |  |  |  |  |  |
|---------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------|
| traumatic skin injuries (wounds, including those after surgical interventions, cracks, scratches) | •                                                                                  | •                                                                                   | •                                                                                   | •                                                                                   | •                                                                                   |
| abscesses (after surgical treatment) caused by microorganisms susceptible to florfenicol          | •                                                                                  | •                                                                                   | •                                                                                   | •                                                                                   | •                                                                                   |
| infectious lesions of horns and hooves                                                            | •                                                                                  | •                                                                                   | •                                                                                   | •                                                                                   |                                                                                     |
| dermatitis of bacterial origin, pyoderma                                                          |                                                                                    |                                                                                     |                                                                                     |                                                                                     | •                                                                                   |
| interdigital dermatitis                                                                           | •                                                                                  | •                                                                                   | •                                                                                   |                                                                                     |                                                                                     |
| acute inflammatory processes around the umbilical cord                                            | •                                                                                  |                                                                                     |                                                                                     |                                                                                     |                                                                                     |
| ulcers, eczema                                                                                    | •                                                                                  | •                                                                                   | •                                                                                   | •                                                                                   | •                                                                                   |
| purulent-necrotic lesions of the extremities                                                      | •                                                                                  | •                                                                                   | •                                                                                   | •                                                                                   | •                                                                                   |




Shake  
the bottle



1-2 sprays



Repeat  
every 12 hours



Course -  
up to 10 days

## IMPORTANT!

Before applying the drug, the affected area must be thoroughly cleaned of exudate and necrotic tissues

**WITHDRAWAL PERIOD:** 0 days. Do not spray the drug on the udders of animals whose milk is used for human consumption or animal feed.

Manufacturer:



www.biotestlab.ua | info@biotestlab.ua

